

Flowsheets

Adult Patient Care Summary

Search (Alt+ Comma)

Hide All Show All

PAIN/COMFORT/SLEEP

Pain/Comfort/Sleep

[REMOVED] Pain Assessment

Pasero Opioid Induced Sedation

Aldrete Postanesthesia

Pain/Comfort/Sleep

COPING/PSYCHOSOCIAL

HEENT

COGNITIVE/NEURO/BEHAVIORAL

NEURO

BEHAVIORAL

RESPIRATORY

CARDIAC

CARDIAC TELEMETRY USE

PERIPHERAL NEUROVASCULAR

GASTROINTESTINAL

GENITOURINARY

ELIMINATION ASSISTANCE

REPRODUCTIVE

BRADEN RISK

SKIN

MUSCULOSKELETAL

NUTRITION

ASPIRATION RISK

NUTRITION

ACC/CS/MONITORING DEVICES

Sign Report

Accordian

Expanded

View All

Admission (Discharged) from 2/20/2020 in SURGIC

	1700	2036	2038
Pain/Comfort/Sleep			
Fever Reduction/Comfort Measures			
Sleep/Rest/Relaxation			
Self Report Pain Scale			
Self report no pain			Numeric PIS
Number Scale Pain Rating 0			
Comfort/Acceptable Pain Level			
Pain Assessment Deferred			
Pain Assessment			Pain Assessment: Numb...
Additional Pain Documentation			
[REMOVED] Pain Assessment: Number Scale (0-10) 02/04/20 0737 back			
Pain Assessment: Number Scale (0-10) -			Pain Body Location
Pain Rating (0-10): Rest			6
Pain Rating (0-10): Activity			6
Frequency			constant
Quality			aching
Pain Radiation to			back
Nonverbal Indicators of Pain			grimace
Factors That Aggravate Pain			movement/positioning
Factors That Relieve Pain			rest
Associated Signs/Symptoms			weakness
Alternative Therapies Interventions			cold application;pillow su...
Pharmaceutical Interventions			around-the-clock dosing ...
Pasero Opioid Induced Sedation Scale			
POSS (Pasero Opioid-Induced Sed Scale)			1 - Awake and alert

Pain Assessment score at 20:38

POSS Score at 20:38

MAR

Documented By: Alvarez Ospina, Melissa

Schedule Date/Time: 02/05/2020 2038

Medication

HYDROMORPHONE (DILAUDID) injection 1 mg : Dose 1 mg : IV Push : EVERY 4 HOURS PRN moderate Pain (PIS 4-6)

Show Flowsheet

Admin Instructions:

If ordered IV Push, administer undiluted at a rate of 0.5 mg over 1 minute. If ordered SC, administer over a few seconds. For SC dose volumes greater than 1.5 mL, use multiple injection sites. RECOMMENDED monitoring for Opiate Naive patients includes BP and Respiratory Monitoring baseline and with pain and level of sedation re-assessment. No BP or Respiratory Monitoring recommended for Maintenance Opiate use patients. RECOMMENDED monitoring for all patients includes Level of Sedation.

Product Instructions:

If ordered IV Push: Administer undiluted over 2 minutes (maximum rate = 0.5 mg/min).

Ordered Admin Amount: 1 mL = 1 mg of 1 mg/mL

Last Admin: 02/05/2020 at 2038 (Given)

Dispense Location: GH Surgery C Pyxis

Frequency: EVERY 4 HOURS PRN

Route: IV Push

Order Dose: 1 mg

Order Questions/Answers

REASON FOR USE: History of chronic pain managed with hydromorphone

Order Start Time: 02/05/2020 at 1500

Order End Time: 02/05/2020 at 1718

PRN Reasons: moderate Pain (PIS 4-6)

References: Micromedex

YNHHS Adult IV Push Guidelines

YNHHS Pediatric and Newborn IVP Guidelines

Linked Line: [REMOVED] Periph IV-single(adult) metacarpal (dorsum of hand), left over-the-needle catheter system 20 gauge [REMOVED] at 2038

Action:

Given

Route: IV Push

Dose: 1 mg

Order Concentration: 1 mg/mL

Scanned Concentration: 1 mg/mL

Pain Medication Administration at 20:38